

Drug Facts

Active ingredient (in each applicator)

Tioconazole 300 mg (6.5%)Vaginal antifungal

Purpose

Use ■ treats vaginal yeast infections

Warnings

For vaginal use only

Do not use if you have never had a vaginal yeast infection diagnosed by a doctor

Ask a doctor before use if you have

- **vaginal itching and discomfort for the first time**
- **lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.**
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to human immunodeficiency virus (HIV) that causes AIDS

When using this product

- do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted disease (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur

Stop use and ask a doctor if

- **symptoms do not get better after 3 days**
- **symptoms last more than 7 days**
- **you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- before using this product read the enclosed brochure and instructions on foil packet for complete directions and information
- adults and children 12 years and over:
 - open the foil packet just before use and remove purple cap
 - insert entire contents of applicator into the vagina at bedtime. Throw applicator away after use.
- children under 12 years of age: ask a doctor

Other information

- this product is a 1-dose treatment, most women do not experience complete relief of their symptoms in just one day. Most women experience some relief within one day and complete relief of symptoms within 7 days.
- if you have questions about vaginal yeast infections, consult your doctor
- store at 20 - 25°C (68 - 77°F)
- see end flap of carton for lot number and expiration date

Inactive ingredients butylated hydroxyanisole, magnesium aluminum silicate, white petrolatum

Questions? call toll-free 1-877-MONISTAT (1-877-666-4782)

← Tear Here →
Do not use if sealed foil packet is torn, open, or incompletely sealed.


MONISTAT[®] 1

1-DAY TREATMENT


TIOCONAZOLE OINTMENT 6.5% VAGINAL ANTIFUNGAL

DIRECTIONS FOR USE

- Tear open foil packet just before using. It is best to use at bedtime.
- Remove applicator and plunger from packet. Applicator is pre-filled with vaginal ointment.
- While firmly holding the capped purple-capped end of the applicator, push the tip of the plunger into the base of applicator.





- ✓ **REMOVE PURPLE CAP** from top of applicator with a pull-twist action.



Insert the applicator

- Lie on your back with your knees bent. Gently insert applicator into the vagina as far as it will go comfortably.
 - Push the plunger into the applicator until it will go no farther. Withdraw the applicator and plunger and dispose of it in the wastebasket. Do not flush.
 - **DO NOT USE TAMPONS** while using this medicine. Use sanitary napkins instead.
- PLEASE READ EDUCATIONAL BROCHURE FOR ADDITIONAL INFORMATION.



Active Made in Italy
Distributed by: INSIGHT Pharmaceuticals, 42600 J5 P2
LLC.
Trevose, PA 19053-8433 25-72-130C
© INSIGHT Pharmaceuticals, LLC. 2014