

MONISTAT® dosing chart for vaginal yeast infections

PRODUCT	PRODUCT DESCRIPTION / DOSING	WHEN TO RECOMMEND
MONISTAT® 1 Day Treatments <i>Maximum Strength</i>	 COMBINATION PACK 1 Ovule® with applicator 1200 mg miconazole nitrate + External itch relief cream 2% miconazole nitrate 	Patient wants <ul style="list-style-type: none"> • Convenience of a single dose • Option to treat during the day or night • Maximum strength treatment • Less messy treatment
	 1 prefilled applicator Tioconazole ointment 6.5% 	<ul style="list-style-type: none"> • Patient has sensitive skin • Patient wants convenience of prefilled product in 1 dose
MONISTAT® 3 Day Treatments <i>Moderate Strength</i>	 COMBINATION PACK 3 Ovules® with applicators 200 mg miconazole nitrate per dose + External itch relief cream 2% miconazole nitrate 	Patient prefers moderate-dose treatment and multiple formulation options (ie, ovules, vaginal suppositories, cream) Recommend for use at bedtime
	 COMBINATION PACK 3 vaginal suppositories with applicators 200 mg miconazole nitrate per dose + External itch relief cream 2% miconazole nitrate 	
	 COMBINATION PACK 3 prefilled applicators (cream)* 200 mg miconazole nitrate per dose + External itch relief cream 2% miconazole nitrate <i>*Can be purchased without external itch relief cream</i> 	
MONISTAT® 7 Day Treatments <i>Low Dose</i>	 COMBINATION PACK Cream with 7 applicators* 100 mg miconazole nitrate per dose + External itch relief cream 2% miconazole nitrate (9 g tube) <i>*Can be purchased without external itch relief cream</i> 	<ul style="list-style-type: none"> • Pregnant patient • Diabetic patient • Patient prefers lower-dose treatment • Patient wants lower-cost treatment Recommend for use at bedtime

Important reminders for your patients

- MONISTAT® may be used during menstrual periods
- Do not have vaginal intercourse while using a MONISTAT® product
- Do not use tampons, douches, spermicides, condoms, or diaphragms while using a MONISTAT® product
- For 3-day and 7-day treatments, doses must be taken over consecutive days for treatment to be effective
- All MONISTAT® yeast infection products can take up to 7 days to fully cure

MONISTAT® provides the broadest-spectrum treatment^{1-3*} plus 4x faster symptom relief^{4†}

Yeast Species	% of Cases	Miconazole (MONISTAT®)	Fluconazole	Butoconazole	Terconazole
<i>C. albicans</i>	70.8%	✓	✓	✓	✓
<i>C. glabrata</i>	18.9%	✓			
<i>C. parapsilosis</i>	5.0%	✓	✓		
<i>C. krusei</i>	2.0%	✓			
<i>S. cerevisiae</i>	1.5%	✓	✓		
<i>C. tropicalis</i>	1.4%	✓	✓		
<i>C. lusitanae</i>	0.2%	✓	✓		
	99.8%				

MONISTAT® relieves yeast infection symptoms

4x faster than fluconazole⁴

MONISTAT® treats the yeast species that cause

99.8% of vaginal yeast infections,¹⁻³ more than fluconazole

>99% of VVC[‡] is caused by 7 yeast species.² MONISTAT® treats them all^{1,3*}

- That's broader coverage than fluconazole—or any other prescription azole^{1,3}
- MONISTAT® treats *C. glabrata*—the most common *non-albicans* species—while prescription azoles do not^{1,2}

MONISTAT® relieves symptoms **4x faster than fluconazole**^{4†}

Provides faster relief from **itching, burning, and irritation**—while curing infections^{4†}

Recommend MONISTAT® to cure your patients' yeast infections and relieve symptoms fast

*vs all first-line prescription drug products

†Based on a clinical study with MONISTAT® 1 Combination Pack Ovule® treatment vs fluconazole

‡VVC=Vulvovaginal candidiasis

References: 1. ISSVD. (2019). Vulvovaginal Candidiasis: Diagnosis and Treatment Tips (Version 1.3) [Mobile application software]. Retrieved from <http://itunes.apple.com>. 2. Richter SS, et al. Antifungal susceptibilities of *Candida* species causing vulvovaginitis and epidemiology of recurrent cases. *J Clin Microbiol.* 2005;43(5):2155-2162. 3. Data on file. Insight Pharmaceuticals LLC, a Prestige Consumer Healthcare company. 4. Phillips N, Bachmann G. Local vs systemic antifungal treatment of yeast vaginitis: comparison of time to symptomatic relief. *J Womens Health Gyn.* 2019;5:1-6.

MONISTAT®